

Prevalence of Tracheostomy and Its Indications in Iran: A Systematic Review and Meta-Analysis

Alireza Alidad ¹, Alireza Aghaz ², Ehsan Hemmati ², Hussein Jadidi ³, Kayvan Aghazadeh ⁴

¹ Speech and Language Pathology, School of Rehabilitation Sciences, Tehran University of Medical Sciences, Tehran, Iran, ² Research Committee and Department of Speech Therapy, School of Rehabilitation Sciences, Isfahan University of Medical Sciences, Isfahan, Iran, ³ Research Committee and Department of Biostatistics and Epidemiology, School of Health, Isfahan University of Medical Sciences, Isfahan, Iran, ⁴Department of ENT, Tehran University of Medical Sciences, Tehran, Iran

Received: 16 December 2018

Accepted: 10 June 2019

Correspondence to: Aghazadeh K

Address: Department of ENT, Tehran University of Medical Sciences, Tehran, Iran

Email address: aghazadeh@sina.tums.ac.ir

Background: Tracheostomy is common among intensive care unit (ICU) patients. Reconsideration of tracheostomy indications in patients can be effective in modifying and reducing irrevocable patient complaints. The present study aimed to analyze the prevalence of tracheostomy indications and to estimate the prevalence of tracheostomy in Iran.

Materials and Methods: In this systematic review and meta-analysis, scientific databases were searched from January 1990 to April 2018. The inclusion criteria were as follows: 1) use of the term “tracheotomy” in the title; and 2) studies conducted in Iran. On the other hand, the exclusion criteria were: 1) studies that did not specify the type of mechanical ventilation; 2) studies that did not quantitatively report the indications; 3) studies without access to the full-text; and 4) case studies, letters to the editor, and/or prefaces. Data were extracted from published reports. Our preliminary results included estimations of tracheostomy indications in Iran.

Results: In the preliminary search, a total of 325 articles were found, 24 of which were considered eligible. Among 2860 patients who had undergone tracheostomy, 21 indications were identified. Decreased mental status, respiratory disease, and tumors were the most frequent indications. The prevalence of tracheostomy was 40.59% in Iran, with the highest and lowest rates reported in Birjand and Ardabil, respectively (136.50 and 6.63 in 100,000 people, respectively) based on the random effects model.

Conclusion: The most prevalent indications in Iran are different from those reported in other countries. This difference may be due to the lack of trained medical personnel and available technologies.

Key words: Tracheostomy, Indications, Prevalence, Systematic Review, Meta-Analysis

INTRODUCTION

Tracheostomy is the formation of a valve or opening into the trachea with the aim of facilitating air passage in upper airway obstructions and improving the discharge of pulmonary secretions or laryngeal storage in patients with long-term intubation (1, 2). The site of incision is usually between the second and third tracheal rings (3). The history of this surgery dates back to pre-Christianity in

Greece (1, 4). According to previous studies, the prevalence of tracheostomy ranges from 6% to 65% (3). In this regard, Fischer et al. in a study from Switzerland showed that 60% of intensive care unit (ICU) patients had undergone tracheostomy in the second week (5), while in a study by Kluge et al. from Germany, the corresponding rate was 90% (6). On the other hand, studies from Iran have reported a prevalence rate of 24% (2).

In the past, the first indication for tracheostomy was the upper airway obstruction due to infectious diseases (7). However, considering the progress in vaccine development and use of antibiotics, infection is not the most prevalent indication for tracheostomy. In most previous studies, avoidance of prolonged intubation is the most common indication for tracheostomy (8, 9). However, the results of some studies are still controversial in this area. For instance, the most prevalent indications in some studies were respiratory diseases, head and neck tumors, and trauma to the jaws and skull, respectively (10). In some other studies, the most common indications were reported to be cardiac and pulmonary diseases, neurological disorders, and airway obstructions (7).

Since tracheostomy is an operation in which the upper airway is bypassed, it may pose serious risks to patients. Also, tracheostomy is one of the most common surgeries in the area of ear, nose, and throat (ENT); therefore, analysis of indications for tracheostomy can be effective in reducing the irreversible complications. The aim of the present study was to analyze the prevalence of tracheostomy indications and to estimate the prevalence of tracheostomy based on the data collected from Iranian studies.

MATERIALS AND METHODS

Search strategy and selection criteria

This systematic review and meta-analysis was conducted according to the PRISMA statement (11). From January 1990 to April 2018, databanks, including the Web of Knowledge, PubMed, Google Scholar, Science Direct, IranDoc, MagIran, and SID, were systematically searched, using the following keywords: “tracheostomy” or “tracheotomy”, “indication”, and “Iran”. The search included the online libraries of all medical universities in Iran. The keyword search in the websites of medical universities was conducted in both Persian and English languages. The Google search engine was also searched to obtain the full-text of the selected articles and to find more information about the subject under study. Moreover, to evaluate the results of keyword search, papers which were

only suggested by the search engines were evaluated for their pertinence to the study.

Two authors conducted separate searches and appraised the title and abstract of each article based on the inclusion criteria. The inclusion criteria were as follows: 1) use of the keyword “tracheotomy” or “tracheostomy” in the title of the article; and 2) performing the study only in Iran. On the other hand, the exclusion criteria were as follows: 1) studies which did not properly define the type of mechanical ventilation (intubation or tracheotomy); 2) studies which did not report the indications in a quantitative manner; 3) studies with unavailable full-texts; and 4) case studies, letters to the editor, and prefaces. In the extracted studies, there was no limitation regarding the type of procedure (surgical or percutaneous) or time of surgery (late or early tracheotomy).

Data extraction and risk of bias assessment

Using a standard form, the two authors (“AA” and “AAD”) separately extracted the data from the studies, including the authors’ names, location of studies, period of studies, number, gender, and average age of participants, and different indications for tracheotomy. Any disagreement between the authors was resolved through discussion with other researchers. The first author of the extracted studies was contacted in case of any ambiguities or to obtain additional information.

Outcomes

The preliminary results of the present study were estimations of all indications for tracheostomy, based on studies on tracheostomy in Iran. Indications reported in each tracheostomy study were assessed. Since these indications varied in the extracted studies, the two authors agreed to identify similar causes, which referred to a specific disorder and to consider them collectively as one indication. The secondary results indicated the prevalence of tracheostomy. Since all studies were conducted during a specific period, the sample size for each study, relative to the population of the city where the study was conducted, could indicate the prevalence of this surgery in that particular city.

The prevalence of tracheostomy in each city was calculated by dividing the number of people undergoing tracheostomy by the city population in the year of its publication. To calculate the overall prevalence of tracheostomy, the total number of tracheostomies reported in the reviewed articles was divided by the total population of cities. The population of each city was determined based on the statistics published by the Statistical Center of Iran.

Statistical analysis

Comprehensive meta-analysis (CMA) software was employed for data analysis. Variance of each study was calculated with regard to binomial distribution, and studies were combined considering the variance and sample size. To calculate the prevalence, 95% confidence interval was used to weigh each of the studies. A random effects model was also applied to combine studies with respect to their heterogeneity. Moreover, Cochran test and I² index were utilized to evaluate the heterogeneity of studies.

Funding source

There was no funding source in this study. Two of the authors had the required access to the necessary information. The decision to publish the article was made by all of the authors.

RESULTS

Selection of studies

Based on the inclusion criteria, a total of 325 articles were found in the preliminary search of domestic and international websites (130 from Google Scholar, 29 from PubMed, 9 from ISI, 2 from Science Direct, 34 from IranDoc, 60 from MagIran, 57 from SID, and 4 from the digital libraries of medical universities in Iran). According to the exclusion criteria, 301 articles were eliminated, including 262 duplicate articles, 33 case reports, and six studies without the full-text manuscript. Finally, 24 full-text articles were included in the meta-analysis. The details of article selection are shown in Figure 1.

Characteristics of studies

Twenty-four studies were published between January 1990 and April 2018. The selected studies were

performed on 2860 patients with tracheostomy in 12 cities of Iran. In 17 articles, the average age of the patients was reported. Based on the findings, the average age of patients with tracheostomy was 49.2 years, and the age range of subjects in all 24 articles was 0-86 years. In 23 articles, 66.1% of the patients were male. These patients underwent tracheostomy due to 21 different indications. The participants' age, gender, and publication date of each study are presented in Table 1.

Frequency of tracheostomy indications

In 24 selected studies, 21 indications were reported. The results of meta-analysis showed that the most common indications for tracheostomy were depressed mental status (19.1%), respiratory disease (14.1%), tumors (10.5%), cardiac problems (9.7%), and laryngeal problems (9.5%). These five indications comprised 62.9% of all indications (Figures 2-6). Depressed mental status (19.1%) was the most prevalent indication, and foreign body was the least common indication for tracheostomy. The frequency and percentage of these 21 indications are presented in Table 2. The obtained results were assessed using the random effects model at 95% confidence interval.

Rate of prevalence

Twenty-four studies reported the prevalence of tracheostomy in 12 cities of Iran. In each study, the rate of prevalence was calculated for each city as the number of tracheostomy cases in every 100,000 people relative to the population of cities. Regarding the considerable heterogeneity of studies ($P < 0.0001$; $I^2 = 100\%$), the prevalence rate was measured for each study according to the random effects model. The prevalence of tracheostomy in the evaluated cities and the overall prevalence of tracheostomy in Iran are shown in Figure 7. The general prevalence of tracheostomy in Iran was 40.59 per 100,000 people (95% CI: 29.95-41.16), with the highest rate reported in Birjand (136.50 in every 100,000 people; 95% CI: 134.91-138.10) and the lowest rate found in Ardabil (6.63 in every 100,000 people; 95% CI: 6.41-6.8) (Figure 7).

Table 1. Trials of tracheostomy Indications meeting inclusion criteria

Indications or Causes of Tracheostomy (Number of Patients)		Characteristics of studies																										
Heart problems	87	Gender (male)	39	-	84	25	46	57	83	233	68	32	125	40	20	58/7	28	14	30	103	10	104	57	24	176	66	165	77
Suicide attempts	6	Age(year)	5/31	-	-	48	20 – 70	65/2	57/3	51/7	55/5	33/48	0 - 86	31/9	58/7	58/4	60/3	0 - 85	64/5	47/8	47/7	-	0 - 85	47/4	53/5	50/8		
Septicemia	84	Sample size (n)	51	402	126	36	80	159	184	360	100	50	131	47	55	16	50	140	12	151	100	32	176	96	210	96		
Hypertension	18	Location (city)	Tehran	Rasht	Tehran	Hamedan	Tehran	Tehran	Kermanshah	Tehran	Tehran	Tabriz	Tehran	Shiraz	Semnan	Tehran	Tehran	Tehran	Tehran	Ahvaz	Zabol	Ardabil	Biljand	Babol	Tabriz	Rasht		
Diabetes	6	Dates	1993 – 2004	2009 - 2011	2002 – 2004	2009- 2010	2013	2012 - 2014	2011 - 2015	2009 - 2014	2009 - 2011	2005 – 2010	1993 – 1999	1999- 2000	2013	1991 - 2000	2012 - 2014	1996 – 2000	2001 - 2010	2001- 2006	2010- 2016	1998- 2001	2000- 2007	2001- 2006	1999- 2004	2010- 2011		
Epilepsy and Tumor in brain	4	Study	Izadi (12)	Saleh (2)	Hasani (13)	Farhanchi (14)	Valandroust (15)	Babaei (16)	Karimpour (17)	Hashemian (18)	Karvandian (19)	Hashemzadeh (20)	Shadab (21)	Hashemi (22)	Hemmati (23)	Hashemzadeh (24)	Ghorbani (25)	Javadi (26)	Bagheri (27)	Nikakhtagh (9)	Malekzadeghan (28)	Amini (29)	Mofateh (30)	Kiakojuri (21)	Toutounchi (32)	Nemati (33)		
Brain injury	161																											
Poisoning	7																											
Airway obstruction	4																											
Blunt/ penetrating neck trauma	25																											
Foreign body	4																											
Laryngeal problems	10																											
Tumor	1																											
Epiglottitis/ supraglottitis	29																											
Maxillofacil fractures or TMJ Ankylosis	53																											
Head and neck surgery	8																											
Pulmonary toilet	50																											
Depressed mental status	241																											
Neuromuscular disease	11																											
Inability to intubate																												
Respiratory disease	12																											

Table 2. Percentage and the frequency of different tracheostomy indications

Indications	Frequency	Percentage*
Depressed mental status	935	%19.1
Respiratory disease	489	%14.1
Tumor	214	%10.5
Heart problems	101	%9.7
Laryngeal problems	271	%9.5
Brain injury	329	%7
Neuromuscular disease	37	%5
Blunt/ penetrating neck trauma	46	%4.5
Septicemia	84	%3.1
Inability to intubate	23	%2.7
Hypertension	18	%2.5
Diabetes	18	%2.3
Epiglottitis/ Supraglottitis	9	%1.7
Head and neck surgery	14	%1.6
Epilepsy and Tumor in brain	11	%1.5
Pulmonary toilet	13	%1.4
Suicide attempts	6	%1.2
Airway obstruction	4	%0.65
Poisoning	2	%0.64
Extensive maxillofacial fractures	2	%0.55
Foreign body	1	%0.55

* This results are reported according to Meta-Analysis




Figure 1. The Flowchart of selected articles




Figure 2. The percentage of tracheostomy cases due to the depressed mental status. The overall percentage of tracheostomy patients due to the depressed mental status in 12 studies was 0.19. The highest percentage was 0.35 in the Hemati study and the lowest rate was 0.05 in the Vatandoust study.




Figure 3. The percentage of tracheostomy cases due to the Respiratory disease. The overall percentage of tracheostomy patients due to the Respiratory disease in 14 studies was 0.14. The highest percentage was 0.23 in the Karvandian study and the lowest rate was 0.01 in the Hashemzadeh study.




Figure 4. The percentage of tracheostomy cases due to the Tumor. The overall percentage of tracheostomy patients due to the Tumor in 7 studies was 0.105. The highest percentage was 0.24 in the Vatandoust study and the lowest rate was 0.05 in the Shadab study.




Figure 5. The percentage of tracheostomy cases due to the Heart problems. The overall percentage of tracheostomy patients due to the Heart problems in 3 studies was 0.097. The highest percentage was 0.195 in the Babae study and the lowest rate was 0.003 in the Malekzadegan study.




Figure 6. The percentage of tracheostomy cases due to the Laryngeal problems. The overall percentage of tracheostomy patients due to the Laryngeal problems in 11 studies was 0.095. The highest percentage was 0.356 in the Hashemzadeh study and the lowest rate was 0.005 in the Malekzadegan study.




Figure 7. The rate of tracheostomy prevalence in every 100 thousand people and the 95% confidence interval in the considered studies based on the model of random effects. The middle point of each line shows the rate of prevalence, and the length of each line indicates the 95% confidence interval in each city. The rhombic sign shows the rate of prevalence combinations in the studies. The overall rate of tracheostomy prevalence in 24 studies was 40.59. The highest prevalence was 136.5 in Birjand and the lowest rate was 6.63 in Ardabil city.

DISCUSSION

The present study was conducted to estimate the indications and prevalence of tracheostomy in a systematic review of studies published in the last three decades in Iran. The results of eligible studies showed that the most common indications for tracheostomy in Iran were different from those reported in other countries. According to the present findings, the most common indications were depressed mental status (19.1%), respiratory diseases (14.1%), tumors (10.5%), cardiac problems (9.7%), laryngeal problems (9.5), and brain injury (7%). On the contrary, in a similar study, the most common indications were cardiopulmonary disease (32%) and neurological disorders (1%) (7).

In another retrospective review, prolonged intubation (35%), upper airway obstruction (28%), neurologic disorders, and craniofacial anomalies (12%) were the most prevalent indications for tracheostomy (34). Differences in the leading cause of tracheostomy are not exclusive to Iran, as every country has its own demographic pattern of health issues, different numbers of trained medical personnel, and different available technologies. In this regard, various reports have been recently published in England, suggesting that the most common indications were infections due to airway obstruction and need for long-term ventilation support (35).

Depressed mental status means loss of consciousness for any reason lead to intubation and artificial ventilator support, which results in tracheotomy if prolonged. Respiratory disease, as the second most common indication for tracheostomy in Iran, includes Subglottic Stenosis, Dysplasia, Pulmonary diseases, Asthma, Pneumonia, Croup, Angina, and Abscess. Tumors, such as Neoplasms of the larynx, thyroid, trachea, and esophagus, are the third most common etiologies in Iran. Brain injury includes Trauma, brain stroke, cerebral hemorrhage, cerebral hypoxia, and multiple traumas. Head and neck surgery includes restoration, cancer, and head and neck surgeries. Moreover, indications related to laryngeal

problems include laryngeal cancer, laryngomalacia, laryngitis, and laryngeal papillomatosis.

The results of the present meta-analysis showed that the general prevalence of tracheostomy in Iran is 40.59 in every 100,000 people. According to this investigation, Birjand has the highest prevalence of tracheostomy among Iranian cities, while Ardabil has the lowest rate. Similar to other meta-analyses, the present study had some limitations. Systematic analyses were mainly performed among cross-sectional studies, and lack of cohort and case-control studies adversely affected the results. Therefore, it is essential to conduct further research in this area. In this study, classification of indications was based on the patients' records before and after surgery. In addition to medical records, the tracheostomy surgeon can select the operation according to the patient's condition. Moreover, two simultaneous diagnoses were suspected in many patients, which could affect the final decision for tracheostomy.

Finally, for estimating the prevalence rate of tracheostomy in Iran, we analyzed the results of studies conducted in 12 different cities according to the city population. In all of these cities, we had no access to the number of people with tracheostomy, and estimations were only based on the published articles. Therefore, regarding the lack of access to the number of people undergoing tracheostomy in these cities and non-consideration of other cities in Iran, the obtained results cannot be confidently generalized to represent all cities in the country, and caution must be taken while interpreting the results.

CONCLUSION

According to studies conducted in the past 28 years in Iran, the most common indications for tracheostomy were decreased mental status, respiratory diseases, and tumors. These findings were not in line with the results of studies performed in other countries. Also, the present review showed that the general prevalence of tracheostomy is 40.59

per 100,000 people in Iran. However, to better understand the epidemiology of tracheostomy in Iran, more extensive studies with a larger sample size are necessary to provide more precise information.

REFERENCES

1. Ganjloo J, Assarroudi A, Rad M. Entering the fractured tracheotomy tube into the left bronchus, A Case Report. *J Sabzevar Uni Med Sci* 2013;20(2):230-235.
2. Saleh S, Molahoseini S, Moghtader O. Study of The effect of tracheostomy care education on anxiety and performance of Family of patients referring to the selected hospital of the Gilan University of Medical Science in 2012. *Sefid Push Sabz J* 2012;38(1):35-44.
3. François B, Clavel M, Desachy A, Puyraud S, Roustan J, Vignon P. Complications of tracheostomy performed in the ICU: subthyroid tracheostomy vs surgical cricothyroidotomy. *Chest* 2003;123(1):151-8.
4. El-Anwar MW, Nofal AA, Shawadfy MA, Maaty A, Khazbak AO. Tracheostomy in the Intensive Care Unit: a University Hospital in a Developing Country Study. *Int Arch Otorhinolaryngol* 2017;21(1):33-37.
5. Fischler L, Erhart S, Kleger G-R, Frutiger A. Prevalence of tracheostomy in ICU patients. A nation-wide survey in Switzerland. *Intensive care med* 2000;26(10):1428-33.
6. Kluge S, Baumann HJ, Maier C, Klose H, Meyer A, Nierhaus A, et al. Tracheostomy in the intensive care unit: a nationwide survey. *Anesth Analg* 2008;107(5):1639-43.
7. Gergin O, Adil EA, Kawai K, Watters K, Moritz E, Rahbar R. Indications of pediatric tracheostomy over the last 30 years: Has anything changed? *Int J Pediatr Otorhinolaryngol* 2016;87:144-7.
8. Yaremchuk K. Regular tracheostomy tube changes to prevent formation of granulation tissue. *Laryngoscope* 2003;113(1):1-10.
9. Nikakhlagh S, Saki N, Mashali L, AKBARI FN, Saleheh F, Joriz M. Survey Of Indication And Early Complications of Tracheostomy In Patients Hospitalized In Ahvaz Imam Khomeini Hospital. *J Med Council Islam Repub Iran* 2013;31(1):45-9.

10. Haspel AC, Coviello VF, Stevens M. Retrospective study of tracheostomy indications and perioperative complications on oral and maxillofacial surgery service. *J Oral Maxillofac Surg* 2012;70(4):890-5.
11. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6(7):e1000097.
12. Izadi F, Delarestaghi M, Pousti S, Kazemi Z, Mir P. A 12-year Survey on the Incidence of Causes and Complications of Pediatric Tracheostomy in Hazrat Rasool-e-Akram Hospital. *Razi Journal of Medical Sciences* 2008;15:55-62.
13. Hassani MD, Zaman B, Motlagh SJ. Comparison between general and local anesthesia in tracheostomies. A two year study in Rasool hospital. *Medical Science Journal of Islamic Azad University-Tehran Medical Branch* 2004;14(1):35-40.
14. Farahanchi A, Sanatkarfar M, Bakhshaie MH, Pouranvari H, Manouchehrian N. The comparison of complications of tracheostomy in percutaneous tracheostomy with surgical tracheostomy. *Avicenna Journal of Clinical Medicine* 2010;17(2):17-23.
15. Vatandoust S, Mohammadi N, Farahani M, Mehran A. The efficacy of education film on quality of life and satisfaction in patient with tracheostomy. *Arak Med Univ J* 2014;17(92):80-7.
16. Babaee T, Alavi SM, Tootoonchi MZ, Farasatkish R, Sadeghpour A, Mohammadzadeh Jooryabi A. The result of modified trans-laryngeal tracheostomy in ventilator-dependent cardiac patients. *Iranian J Anesthesiology and Critical Care*. 1995.
17. Karimpour HA, Vafaii K, Chalechale M, Mohammadi S, Kaviannezhad R. Percutaneous Dilatational Tracheostomy via Griggs Technique. *Arch Iran Med* 2017;20(1):49-54.
18. Hashemian SM, Digaleh H; Massih Daneshvari Hospital Group. A Prospective Randomized Study Comparing Mini-surgical Percutaneous Dilatational Tracheostomy With Surgical and Classical Percutaneous Tracheostomy: A New Method Beyond Contraindications. *Medicine (Baltimore)* 2015;94(47):e2015.
19. Karvandian K, Yousefian M, Khan ZH, Baigmohammadi T, Shabani S. Comparative clinical trial between Ciaglia and Griggs techniques during tracheostomy performed in patients admitted to intensive care unit. *Acta Med Iran* 2012;50(8):525-9.
20. Hashemzadeh S, Hashemzadeh K, Kakaei F, Aligholipour R, Ghabili K. Surgical treatment of postintubation tracheal stenosis: Iranian experience of effect of previous tracheostomy. *Int J Gen Med* 2012;5:93-8.
21. Shadab H. Investigating the Causes of Tracheostomy in Amiralam Hospital and Oral Surgery Department of Shariati Hospital (from 1992 to 1998). dental College. Tehran: Tehran Uni Med Sci 2000. [Thesis]
22. Hashemi SB, Gamdomi B, Derakhshandh V. Evaluation of the causes of upper airway obstruction leading to tracheostomy in Shiraz. *Iranian J Otorhinolaryngol* 2006;18(43):5-8.
23. Hemmati H, Forozeshfard M, Hosseinzadeh B, Hemmati S, Mirmohammadkhani M, Bandari R. Tracheostomy in Patients Who Need Mechanical Ventilation: Early or Late? Surgical or Percutaneous? A Prospective Study in Iran. *Indian J Surg* 2017;79(5):406-411.
24. Hashemzadeh S, Banazadeh M, Kakaei F. Mediastinal tracheostomy in Tehran Imam Khomeini's hospital from january 1991 till december 2000. *Acta Medica Iranica* 2006;44(4):268-72.
25. Ghorbani-Abdehghah A, Mirzaei-Baboli K, Ashouri M, Tajedin A, Gohari-Moghadam K. Comparison of Standard and Percutaneous Tracheostomy Complications in Shariati Hospital in Iran. *Academic Journal of Surgery* 2015;2(1-2):7-10.
26. Javadi M, Arbabi M, Sedghi M, Haghdad A. The study of Indications and Complications of tracheotomy in Hazrat Rasool-e-Akram hospital from 1996-2000. *Razi Journal of Medical Sciences* 2003;10(36):541-5.
27. Bagheri R, Rahim MB, Majidi M, Tabari A, Akhlaghi S, Tabari A. Anterior mediastinal tracheostomy for malignancy: analysis of 12 cases. *Asian Cardiovasc Thorac Ann* 2013;21(2):187-92.
28. Malekzadegan A, Kamkar P, Sargazi A. Evaluation of Causes and Early Complications of Tracheostomy in the Patients Admitted to ICU of Amir-al-Momenin Hospital of Zabol from 2010-2016. *Iran J Surg* 2016;24(3):53-60.
29. Amini y. Prevalence and causes of tracheostomy in 1998-2001 in Fatemi and Alavi hospitals of Ardabil. Medical College. Ardabil: Ardabil UniMed sci 2001. [Thesis]

30. Mofateh M, Golboei Mosavi S. Epidemiology of tracheotomy cases in Birjand, s Hospitals during 2000-2007. *J Birjand Uni Med Sci* 2010;17(3):221-27.
31. Kiakojouri K, Amiri P, Ahmadi H, Madadian M. Indication and Early Complications of Tracheostomy in the Intensive Care Unit Patients in Shahid Beheshti and Shahid Yahyanejad Hospital (Babol, Iran 2001-2006). *Journal of Babol University of Medical Sciences* 2009;11(1):67-71.
32. Toutounchi S, Doust MN, Bagheri RKA. Survey of the Complications of tracheotomy in Tabriz Imam Khomeini hospital during 5 years (2000-2005). *Med J Tabriz Uni Med Sci* 2008;29(4):31-36.
33. Nemati S, Banan R, Kazem Nezhad E, Movahedi H, Bakhshi F, Taravat F. Evaluation of post-tracheostomy care status in hospitals of rasht. *Journal of Guilan University of Medical Sciences* 2013;22(86):32-9.
34. Lawrason A, Kavanagh K. Pediatric tracheotomy: are the indications changing? *Int J Pediatr Otorhinolaryngol* 2013;77(6):922-5.
35. Douglas CM, Poole-Cowley J, Morrissey S, Kubba H, Clement WA, Wynne D. Paediatric tracheostomy-An 11 year experience at a Scottish paediatric tertiary referral centre. *Int J Pediatr Otorhinolaryngol* 2015;79(10):1673-6.